

# Exogenous Hydrogen Sulfide Ameliorates Seed Germination and Seedling Growth of Cauliflower Under Lead Stress and Its Antioxidant Role

Zhen Chen<sup>1</sup> · Beifen Yang<sup>1</sup> · ZhiKui Hao<sup>2</sup> · Jiaqian Zhu<sup>1</sup> · Yan Zhang<sup>1</sup> · Tiantian Xu<sup>1</sup>

Received: 17 September 2016 / Accepted: 20 March 2017 / Published online: 12 April 2017  
© Springer Science+Business Media New York 2017

**Abstract** Lead (Pb) is a widespread ecosystem pollutant and affects food security and public health. Hydrogen sulfide (H<sub>2</sub>S) plays prominent roles in mediating a variety of responses to stresses. The effects of sodium hydrosulfide (NaHS), a fast releaser of H<sub>2</sub>S, on cauliflower (*Brassica oleracea* L. var *botrytis* L. cv. Xiahua 60 d) seed germination and seedling growth under lead acetate stress were investigated in the present study. Pb (0.25 and 0.5 mM) stresses markedly inhibited seed germination and seedling growth, whereas the inhibition was effectively mitigated by NaHS application. Germination percentage, root length, shoot length, and fresh weight of single seedling significantly increased. In addition, NaHS elevated endogenous H<sub>2</sub>S contents and reduced malonyldialdehyde, superoxide anion (O<sub>2</sub><sup>·-</sup>), and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) production, thereby preventing oxidative damage from Pb or Pb and antioxidant enzyme inhibitor (diethyldithiocarbamate or 3-amino-1,2,4-triazole) dual stresses. The protective roles of NaHS were equivalent to the ROS scavengers, 4,5-dihydroxy-1,3-benzene disulfonic acid<sup>?</sup> and *N,N*-Dimethylthiourea. Moreover, NaHS elevated non-protein thiols and total glutathione levels to chelate Pb or scavenge ROS directly. Our results demonstrated the strong protective and antioxidant roles of H<sub>2</sub>S.

**Keywords** Pb stress · H<sub>2</sub>S · Cauliflower · ROS scavenging · Antioxidant effect

## Introduction

Lead (Pb), a ubiquitous industrial and environmental pollutant, has been listed by the Agency for Toxic Substances and Disease Registry (ATSDR 2005) as a major heavy metal pollutant (Sidhu and others 2016a). It is easily absorbed by crops through contaminated air, water, and soil, and then threatens food safety and public health, particularly children and infants (Dewanjee and others 2015; Kim and others 2016; Wang and others 2015). Lead acetate (PbAc) is known to induce kidney damage, hepatotoxicity, neurotoxicity, oxidative imbalance, fibrosis, apoptosis, and reproductive system diseases (Abdel Moneim 2016; Abdou and Hassan 2014; Mahmoud and Sayed 2016; Rao and others 2016). In plants, Pb exposure also exerts deleterious effects on a series of physiological processes including seed germination, root elongation, cell division, photosynthesis, respiration, nutrients uptake and hormonal equilibrium, therefore affecting morphology, growth, and yield (Kanwal and others 2014; Lamhamdi and others 2011, 2013; Sharma and Dubey 2005; Wierzbicka and Obidzińska 1998; Yang and others 2016). Ultrastructural studies have identified that stomatal size and conductance were reduced, thylakoid membranes dissolved, starch grains vanished, cells shrunk, mitochondria and endoplasmic reticulum (ER) were destroyed, the structure of the nucleolus changed, and membrane permeability was modified in plant leaf and root cells under Pb toxicity (Ali and others 2014; Jiang and others 2014; Khan and others 2016b). High concentrations of Pb alter various gene expressions in different tissues (Kovalchuk and others 2005; Li and others 2015; Liu and

✉ Zhen Chen  
jennie1980@163.com; chenzh@tzc.edu.cn

others 2009; Reis and others 2015). Further, excess Pb in plant cells induces molecular damage mainly through triggering bursts of reactive oxygen species (ROS) including superoxide anion ( $O_2^{\cdot-}$ ), hydrogen peroxide ( $H_2O_2$ ), and hydroxyl radical ( $\cdot OH$ ) (Hu and others 2015). These excessive ROS lead to irreversible peroxidation of lipids, proteins, photosynthetic pigments, RNA, and DNA, thereby affecting cell structure and viability (Verma and Dubey 2003). Plants have developed efficient antioxidant systems, including enzymatic and non-enzymatic antioxidants, to combat the toxicity of ROS and oxidative stress (Mittler 2002). Superoxide dismutase (SOD), as the first defense line, converts  $O_2^{\cdot-}$  into  $H_2O_2$ , and then  $H_2O_2$  can be quickly decomposed into  $O_2$  and  $H_2O$  by catalase (CAT), peroxidase (POD), and ascorbate peroxidase (APX). Another  $H_2O_2$  detoxification way is the Halliwell–Asada cycle with the participation of the antioxidants—ascorbate (AsA), glutathione (GSH), and glutathione reductase (GR) (Chen and others 2013b). Sustainable agriculture has emerged as a chief goal in the face of changing environmental conditions. So efficient approaches for detoxification of Pb are greatly needed. Some low cost and viable exogenous substances, such as salicylic acid (SA), abscisic acid (ABA), nitric oxide (NO), flavonoid preparation, ethylene diamine tetraacetic acid (EDTA), GSH, 20-hydroxyecdysone, and organic chelants have been used to assist in the detoxification for Pb stress (Chen and others 2007; Kanwal and others 2014; Kaur and others 2015; Khan and others 2016a, b; López-Orenes and others 2014; Lamhamdi and others 2016; Zhao and others 2009).

Just like NO, the new gaseous signal molecule hydrogen sulfide ( $H_2S$ ) has been recommended for mediating a variety of physiological processes and responses to abiotic stresses (Chen and others 2011a; Fang and others 2014; Christou and others 2013; Hu and others 2012; Li and others 2013; Shi and others 2013; Wang and others 2012; Zhang and others 2011).  $H_2S$  effectively alleviated metal toxicity from copper (Cu), aluminum (Al), cadmium (Cd), and zinc (Zn) (Chen and others 2013a; Liu and others 2016; Mostofa and others 2015; Zhang and others 2008). The protective roles of  $H_2S$  against Pb stress were attributed to the elevated activities of SOD, POD, CAT, and APX and the decreased influx and transport of Pb in cotton (Bharwana and others 2014). So the decreases in electric conductivity (EC), malonyldialdehyde (MDA), and  $H_2O_2$  contents demonstrated lowered oxidative damage by  $H_2S$  supplementation. Exogenous  $H_2S$  also had promotive effects on plant growth, root morphology, chlorophyll contents, net photosynthesis rate, and ultrastructural changes, such as integrated chloroplasts with well-developed thylakoid membranes and starch grains in mesophyll cells, and mature mitochondria, golgi bodies, and long ER in the root tip cells of *Brassica napus* (Ali and others 2014).

Moreover,  $H_2S$  alleviated the hazardous impact of elements on plants by increasing cell wall-related pectin methylesterase (PME), expansions, citrate, or plasma membrane (PM)  $H^+$ -ATPase (Chen and others 2013a; Wang and others 2010). On the other hand,  $H_2S$  might serve as a prominent signal transmitter to activate downstream signal transductions and therefore attenuate the stresses from heavy metals (Li and others 2012a; Shi and others 2014). However, systematically understanding the protective mechanism of  $H_2S$  is quite complicated and challenging.

Cauliflower, belonging to Brassicaceae, is a popular vegetable grown throughout the world for its abundant proteins, minerals, vitamins, and metabolites which protect mankind from heart diseases and certain cancers (Keck and Finley 2004). Plant species related to Brassicaceae are generally considered as sensitive indicators or phytoremediators for their fast growth, higher biomass, and adsorptive ability of heavy metals (Ali and others 2014; Gupta and Gupta 2015; Sidhu and others 2016). However, the role of  $H_2S$  in alleviating Pb stress is still unknown at least when it comes to cauliflower. In this study, sodium hydrosulfide (NaHS), a common donor of  $H_2S$ , was applied to study the effects of exogenous  $H_2S$  treatments on seed germination and seedling growth of cauliflower under Pb stresses and the results would help understanding of the antioxidant effects of  $H_2S$ .

## Materials and Methods

### Plant Culture and Treatment

Seeds of cauliflower (*Brassica oleracea* L. var *botrytis* L.) cv. Xiahua 60 days were surface sterilized in 15% (v/v) sodium hypochlorite (NaClO) solution for 10 min followed by thorough rinsing. After being soaked for 3 h, every 20 seeds were cultivated on sterilized solid medium (1% agar) with 0, 0.1, 0.2, 0.3, 0.6, 0.9, 1.5, 3.0 mM NaHS in transparent glass bottles in duplicate for 12 h. Then the seeds were divided into three groups and treated by 0, 0.25, or 0.5 mM lead acetate [ $Pb(CH_3COO)_2 \cdot 3H_2O/PbAc$ ] on fresh media individually. Seeds were placed in the dark at  $(25 \pm 1)^\circ C$  for 3 days and then cultured under cycles of 14 h light/10 h dark photoperiod with 80–100  $\mu mol/(m^2 \cdot s)$  illumination intensity for another 4 days. The phenotype was observed and germination percentage (GP) and biomass were measured.

For the physiological mechanism, cauliflower seedlings were cultured in half-strength Hoagland's solution for 20 days (Hoagland and Arnon 1950). The culture solutions were refreshed every 4 days. After that, the uniform seedlings were treated with 0 or 0.5 mM PbAc for 24 h, then transferred to the culture solutions added with NaHS, DDC

(diethyldithiocarbamate, a Cu/Zn-SOD inhibitor), Tiron (4,5-dihydroxy-1,3-benzene disulfonic acid, a  $O_2^{\cdot-}$  scavenger), AT (3-amino-1,2,4-triazole, a catalase inhibitor), or DMTU (*N,N*-dimethylthiourea, a  $H_2O_2$  scavenger) for another 48 h to prevent direct precipitation of Pb. There were ten treatment combinations for enzymatic antioxidant assays: (1) CK, control without any reagents for treatment; (2) 0.2 mM NaHS; (3) 0.5 mM PbAc; (4) 0.5 mM PbAc + 0.2 mM NaHS; (5) 0.5 mM PbAc + 0.3 mM DDC; (6) 0.5 mM PbAc + 0.3 mM DDC + 0.2 mM NaHS; (7) 0.5 mM PbAc + 1.0 mM Trion; (8) 0.5 mM PbAc + 2  $\mu$ M AT; (9) 0.5 mM PbAc + 2  $\mu$ M AT + 0.2 mM NaHS; (10) 0.5 mM PbAc + 1 mM DMTU. Another 6 treatment combinations were designed for non-enzymatic antioxidant assays: (1) CK; (2) 0.2 mM NaHS; (3) 0.3 mM NaHS; (4) 0.5 mM PbAc; (5) 0.5 mM PbAc + 0.2 mM NaHS; (6) 0.5 mM PbAc + 0.3 mM NaHS. Leaves were used for physiological and biochemical parameter measurements.

### Quantification of Leaf $H_2S$

$H_2S$  contents in leaves were detected according to the previously described procedure (Christou and others 2013). Leaf samples (0.3 g) were grounded into powder with liquid nitrogen and homogenized in 1 ml potassium phosphate buffer (100 mM, pH 7.0) with 10 mM EDTA. After centrifugation at  $12,000\times g$  for 15 min at  $4^\circ C$ , supernatants were collected. The initial absorbance at 412 nm was recorded with Multiskan Spectrum (Thermo, USA). Then 200  $\mu$ l extraction was mixed with 4  $\mu$ l of 20 mM 5,5-dithiobis-2-nitrobenzoic acid (DTNB) and absorbance at 412 nm was measured every 5 min. The steady values of 10 min were chosen for quantification and  $H_2S$  concentration was calculated based on a standard curve of NaHS reagent.

### MDA and ROS Determinations

The concentrations of  $O_2^{\cdot-}$  and  $H_2O_2$  in cauliflower leaf samples were measured according to the regular methods (Jana 1981; Mishra and Singhal 1991). MDA is a byproduct of lipid peroxidation and its level was determined following the method described by Wang and others (2009) with small modifications. Fresh leaves (0.5 g) were ground with 5 ml 10% (w/v) trichloroacetic acid (TCA). After centrifugation at  $8000\times g$  for 10 min, 2 ml of the supernatant was collected and mixed with 2 ml 0.6% TBA solution (dissolved in 10% TCA). Mixtures were kept in a boiling-water bath for 15 min, and then immediately cooled and centrifuged at  $8000\times g$  for 10 min. Then the absorbances of the final supernatants at 450, 532, and 600 nm were recorded. The MDA contents were calculated as follows: ( $\mu$ M MDA) =  $6.45(A_{532} - A_{600}) - 0.56A_{450}$ , and the final contents were presented as nmol/g fresh weight (Fw).

### Measurements of Antioxidant Enzyme Activities

Cauliflower leaves (0.5 g) were homogenized in 3 ml ice-cold 50 mM potassium phosphate (pH 7.8) buffer containing 1% polyvinyl pyrrolidone (PVP, w/v) for enzyme extractions. The homogenates were centrifuged at  $8000\times g$  and  $4^\circ C$  for 20 min and the supernatants were collected. Superoxide dismutase (SOD, EC 1.15.1.1) activity was measured by its ability to inhibit photochemical reduction of nitroblue tetrazolium chloride (NBT) at 560 nm (Beauchamp and Fridovich 1971). One unit of SOD was defined as the amount of enzyme causing 50% inhibition of NBT reduction under light for 6 min. Catalase activity (CAT, EC 1.11.1.6) was assayed according to the reported method (Aebi 1983). The decrease of absorbance at 240 nm was recorded and the final CAT activity was estimated using the extinction coefficient of 39.4 l/(mmol-cm). One unit of enzyme activity was defined as the reduction of 1 nM  $H_2O_2$  per minute. Peroxidase (POD, EC 1.11.1.7) activity was determined following the procedure described by Rao and others (1997). The increase of absorbance at 470 nm was read continuously for 2 min and the extinction coefficient was 26.6 l/(mmol-cm). One unit of POD activity was defined as the formation of 1 nM tetraguaiacol per minute. The activity of ascorbate peroxidase (APX, EC 1.11.1.11) was measured according to the previously published method (Nakano and Asada 1981). Three ml of the reaction mixture consisted of 50 mM potassium phosphate buffer (pH 7.0), 0.1 mM  $Na_2$ -EDTA, 0.5 mM AsA, 0.1 mM  $H_2O_2$ , and 300  $\mu$ l enzyme extract. The decrease of absorbance at 290 nm was recorded and the extinction coefficient was 2.8 l/(mmol-cm). One unit of APX activity was defined as the oxidation of 1 nM ascorbate per minute. Glutathione reductase (GR, EC 1.6.4.2) activity was assayed in terms of the method described by Schaedle and Bassham (1977). The reaction mixture (2.5 ml) contained 100 mM Tris-HCl buffer (pH 7.8), 2 mM EDTA, 50  $\mu$ M NADPH, 0.5 mM glutathione disulfide (GSSG), and 100  $\mu$ l enzyme extract. The extinction coefficient for GR activity calculation was 6.22 l/(mmol-cm) and one unit of GR activity was defined as the reduction of 1 nM NADPH at 340 nm per minute.

### Quantification of Non-protein Thiols (NPT) and Glutathione (GSH)

Leaf samples (0.2 g) were ground in 2 ml 5% (w/v) sulfosalicylic acid solution containing 6.3 mM diethylenetriamine-pentaacetic acid (DPTA) under an ice bath. The mixture was centrifuged at  $4^\circ C$  under  $10,000\times g$  for 10 min, and then the supernatant was immediately collected. Fifty micro liters of the supernatant was mixed with 100  $\mu$ l 0.5 M  $K_2HPO_4$  solution. Two minutes later, the initial absorbance at 412 nm was recorded. Then 5  $\mu$ l DTNB solution (6 mM DTNB

dissolved in 100 mM phosphate buffer, pH 7.6, with 5 mM EDTA) was supplemented. Ten minutes later, absorbance at 412 nm was measured again. The steadily increased absorbance was used for calculating NPT concentration with an extinction coefficient of 13.6 l/(mmol·cm) (Mendoza-Cozatl and others 2008). Total GSH was assayed using an ELISA kit (S0053) from Beyotime Institute of Biotechnology (China). The assays were performed according to the manufacturer's manual. The standard calibration curve was plotted with 0.5–15  $\mu$ M GSSG (Chen and others 2012).

### Statistical Analysis

Values were expressed as means  $\pm$  standard deviations (SD) of at least triplicate independent experiments and measurements. Significant differences analyses were performed

with Student *t* test for pairs or one-way analysis of variance for groups (Duncan's new multiple range test).

## Results

### Exogenous H<sub>2</sub>S Promoted Seed Germination and Seedling Growth of Cauliflower Under Pb Stresses

Under normal conditions, the germination percentage (GP), root length (RL), shoot length (SL), and fresh weight of a single seedling (SW) of 7-day-old seedlings of cauliflower 'Xiahua 60 d' were 90.0%, 6.21, 2.06 cm, and 0.0737 g, respectively (Fig. 1a–d). NaHS (0.1–0.9 mM, an H<sub>2</sub>S fast releaser) preteigheffspengsnle seedlinr growte


(Fig. 2a–d). The results showed that the germination percentage (GP) of cauliflower seedlings under Pb stress (100 mg/L) was significantly lower than that of the control (90.0%) (Fig. 2a). The addition of NaHS (0.1–0.9 mM) significantly increased the GP of cauliflower seedlings under Pb stress (100 mg/L) (Fig. 2a). The results showed that the germination percentage (GP) of cauliflower seedlings under Pb stress (100 mg/L) was significantly lower than that of the control (90.0%) (Fig. 2a). The addition of NaHS (0.1–0.9 mM) significantly increased the GP of cauliflower seedlings under Pb stress (100 mg/L) (Fig. 2a).

compared to the control level (Fig. 1b). Three mM NaHS also reduced SL and SW (Fig. 1c, d), indicating that higher concentrations of NaHS might be toxic to cauliflower seeds.

Pb (0.25 and 0.5 mM) significantly inhibited cauliflower seedling growth after seed germination (Fig. 1). RL was only 1.97 or 0.08 cm, with sharp decreases of 68.28 or 87.12% in comparison with the control. Meanwhile, GP, SL, and SW declined by 7.41 or 18.56%, 37.38 or 54.37%, and 11.80 or 43.01%, respectively. NaHS pretreatment at concentrations of 0.1–0.9 mM effectively promoted seed germination and seedling growth under Pb stresses whereas higher concentrations of 1.5–3.0 mM had no significant effect. NaHS (0.2 mM) showed the optimum effect, and GP, RL, SL, and SW increased by 20.00 or 27.27%, 31.42 or 62.09%, 30.85 or 32.16%, 17.00 or 36.21%, respectively, compared to the condition of 0.25 or 0.5 mM Pb alone.

### Leaf H<sub>2</sub>S Contents

Under normal conditions, H<sub>2</sub>S in leaves was kept at a low level of about 6.66 nmol/g Fw (Fig. 2). Exogenous root application of NaHS led to a significant burst of H<sub>2</sub>S in leaves, up to approximately 30 nmol/g Fw. Pb exposure also induced some H<sub>2</sub>S production and the concent of H<sub>2</sub>S




**Fig. 2** Effects of 0.2 mM NaHS root treatment on leaf H<sub>2</sub>S contents. (1) CK, control; (2) NaHS, 0.2 mM NaHS treated for 48 h; (3) Pb, 0.5 mM PbAc treated for 24 h; (4) Pb+NaHS, 0.5 mM PbAc treated for 24 h then transferred to 0.2 mM NaHS treatment for 48 h; (5) Pb+DDC, 0.5 mM PbAc treated for 24 h then transferred to 0.3 mM DDC treatment for 48 h; (6) Pb+DDC+NaHS, 0.5 mM PbAc treated for 24 h then transferred to 0.3 mM DDC and 0.2 mM NaHS treatment for 48 h; (7) Pb+Tiron, 0.5 mM PbAc treated for 24 h then transferred to 1.0 mM Tiron treatment for 48 h; (8) Pb+AT, 0.5 mM PbAc treated for 24 h then transferred to 2 μM AT treatment for 48 h; (9) Pb+AT+NaHS, 0.5 mM PbAc treated for 24 h then transferred to 2 μM AT and 0.2 mM NaHS treatment for 48 h; (10) Pb+DMTU, 0.5 mM PbAc treated for 24 h then transferred to 1 mM DMTU treatment for 48 h, the same as follows. Data are presented as the mean ± SD (n=4). Values with *different lowercase* mean the significant difference at P<0.05 level

was as much as 2.11-fold of the control, but much lower than NaHS treatment alone. The O<sub>2</sub><sup>·-</sup> scavenger, Tiron, and antioxidant enzyme inhibitors, DDC and AT, had no obvious effects on the changes of H<sub>2</sub>S content in seedlings after Pb stress. The H<sub>2</sub>O<sub>2</sub> scavenger, DMTU, treated after Pb stress, significantly enhanced H<sub>2</sub>S level, compared to Pb treatment alone. When exogenous NaHS was supplemented to the solutions of Pb or Pb and antioxidant enzyme inhibitor co-treatments, H<sub>2</sub>S contents in leaves were all greater than 30 nmol/g Fw (Fig. 2), proving the effective release of H<sub>2</sub>S.

### Exogenous H<sub>2</sub>S Alleviated Oxidative Stress Caused by Pb Stress


Exposure to 0.5 mM Pb caused severe oxidative stress to cauliflower seedlings. MDA, O<sub>2</sub><sup>·-</sup>, and H<sub>2</sub>O<sub>2</sub> contents in leaves under Pb stress rose up to 2.04-, 2.39-, and 2.55-fold of controls (Figs. 3, 4). NaHS alone did not induce excessive MDA and O<sub>2</sub><sup>·-</sup> production but slightly increased H<sub>2</sub>O<sub>2</sub> content. However, the generation of ROS and MDA triggered by Pb was markedly diminished by NaHS treatment (Figs. 3, 4). The low oxidative damage proved the protective effect of H<sub>2</sub>S.

For further investigations, antioxidant enzyme inhibitors and ROS scavengers were used in this study. When SOD activity was inhibited by DDC, O<sub>2</sub><sup>·-</sup> could not be effectively converted into H<sub>2</sub>O<sub>2</sub> and its production rate remained at a high level (Figs. 4, 5a). When NaHS was supplemented to this condition of Pb+DDC, the O<sub>2</sub><sup>·-</sup> production rate decreased sharply to the level when O<sub>2</sub><sup>·-</sup> was eliminated by its scavenger, Tiron (Fig. 4). In addition, when CAT activity was inhibited by AT, H<sub>2</sub>O<sub>2</sub> decomposition was partly



**Fig. 3** Effects of H<sub>2</sub>S, antioxidant enzyme inhibitors, and ROS scavengers on malondialdehyde (MDA) contents in the leaves of cauliflower seedlings under Pb stress. Data are presented as the mean ± SD (n=4). Values with *different lowercase* mean the significant difference at P<0.05 level

blocked. NaHS supplement reduced the H<sub>2</sub>O



**Fig. 5** Activity analysis of antioxidant enzymes in the leaves of cauliflower seedlings under Pb stress, antioxidant enzyme inhibitors, and ROS scavenger treatments. **a** SOD; **b** CAT; **c** POD; **d** APX; **e** GR.

Data are presented as the mean ± SD (*n*=4). Values with *di* *erent lowercase* mean the significant difference at *P*<0.05 level

**Table 1** Effects of exogenous H<sub>2</sub>S on NPT and total GSH contents in cauliflower leaves under Pb stress

| Treatments                          | NPT (nmol/g Fw) | Total GSH (nmol/g Fw) |
|-------------------------------------|-----------------|-----------------------|
| CK                                  | 17.90 ± 1.22d   | 26.16 ± 5.93c         |
| 0.2 mM H <sub>2</sub> S             | 38.47 ± 2.04c   | 30.85 ± 5.03c         |
| 0.3 mM H <sub>2</sub> S             | 22.96 ± 3.89d   | 28.42 ± 6.20c         |
| 0.5 mM Pb                           | 137.68 ± 11.44b | 64.88 ± 5.84b         |
| 0.5 mM Pb + 0.2 mM H <sub>2</sub> S | 174.22 ± 10.95a | 81.58 ± 6.24a         |
| 0.5 mM Pb + 0.3 mM H <sub>2</sub> S | 168.76 ± 16.18a | 81.10 ± 8.60a         |

Data are presented as the mean ± SD (*n*=4). Values with *di* *erent lowercase* mean the significant difference at *P*<0.05 level

that low concentrations of NaHS had slight impacts on cauliflower whereas higher concentrations of 1.5–3.0 mM significantly inhibited seedling growth. It is consistent with the findings from Chen and others (2013a). So the effects of H<sub>2</sub>S is concentration dependent. NaHS at 0.2 mM had the greatest effect on Pb detoxification in cauliflower.

Yang and others (2016) reported that PbAc inhibited six forest tree seed germination and the inhibitory effect order from strong to weak was RL, SL, GP, and SW. Similar results were also found in our study (Fig. 1). Roots are directly exposed in the soil or solution and accumulated the highest amounts of Pb compared to shoots or leaves (Ali and others 2014; Bharwana and others 2014; Hu and others 2015; Sidhu and others 2016). On the other hand, the root endodermis might act as a barrier to the absorption of Pb into the stele and transport to shoots (Reis and others 2015). When Pb entered the radicle cells, it bound

with nucleic acid, disrupted mitosis, and then influenced cell division and root elongation (Yang and others 2016). However, exogenous H<sub>2</sub>S distinctly lowered the Pb concentration in roots and shoots (Ali and others 2014; Bharwana and others 2014). But the mechanism for preventing Pb uptake by intracellular H<sub>2</sub>S is still unclear. The improved effect of H<sub>2</sub>S on root morphology and cell structure, that is, a well-developed nucleus with continuous cell membrane, has been proven by Ali and others (2014) and this may be the crucial contribution to the enhanced root growth.

PbAc exposure significantly up-regulated ROS production both in vivo and in vitro (Dewanjee and others 2015; Dua and others 2016). In this investigation, the decreases in MDA, O<sub>2</sub><sup>-</sup>, and H<sub>2</sub>O<sub>2</sub> contents confirm the protective role of H<sub>2</sub>S against Pb stress in cauliflower (Figs. 3, 4). It is in line with the findings in cotton and tall fescue under Pb stress (Bharwana and others 2014; Hu and others 2015). Similar results about H<sub>2</sub>S were also found in plants under other heavy metal stresses (Chen and others 2013a; Mostofa and others 2015; Zhang and others 2008). Reduced oxidative damage under stresses was usually attributed to enhanced antioxidant enzyme activities stimulated by H<sub>2</sub>S application (Bharwana and others 2014; Li and others 2014; Mostofa and others 2015; Sun and others 2013; Zhang and others 2010). In contrast, it should be noted that the activities of SOD, CAT, POD, APX, and GR were not up-regulated by H<sub>2</sub>S in this study (Fig. 5). The same phenomenon was also found by Chen and others (2013a), who described that in plants treated with 200 μM H<sub>2</sub>S pre-application plus Al stress, the decreased SOD, APX, and POD activities matched the lower H<sub>2</sub>O<sub>2</sub> level and reduced oxidative damage. The authors considered that H<sub>2</sub>S diminished the need for antioxidant enzymes by keeping the ROS levels low in plant tissues. But there were no explanations about how this happened in plants. In the bacteria of *Shewanella oneidensis*, Wu and others (2015) reported that H<sub>2</sub>S had no influence on SOD activity but depressed CAT and POD activities, suggesting that neither SOD nor CAT



still needed for comprehensive elucidation of the protective mechanisms from H<sub>2</sub>S.

In conclusion, 0.25 and 0.5 mM Pb stress significantly inhibited seed germination and seedling growth of cauliflower. However, applications of exogenous H<sub>2</sub>S (NaHS) with suitable concentrations markedly increased leaf H<sub>2</sub>S content and alleviated Pb phytotoxicity. Germination percentage, root length, shoot length, and fresh weight notably increased whereas MDA, O<sub>2</sub><sup>·-</sup>, and H<sub>2</sub>O<sub>2</sub> production declined. H<sub>2</sub>S prevented oxidative damage from Pb or Pb and antioxidant enzyme inhibitor (DDC or AT) dual stresses, and the protective role of H<sub>2</sub>S was parallel to the ROS scavengers, Tiron, and DMTU. Moreover, H<sub>2</sub>S elevated NPT and total GSH levels to chelate Pb or scavenge ROS directly. Our results demonstrated the strong protective and antioxidant role of H<sub>2</sub>S.

**Acknowledgements** This work was supported by Taizhou University Scientific Research Project (2014PY024), Zhejiang Provincial Key Laboratory of Plant Evolutionary Ecology and Conservation open projects (EEC2014-08), and Taizhou Municipal Innovation Team and Key Laboratory Independently Design and Technology Projects (1403ky06).

**Author Contribution** ZC experiment design, result analysis and discussion, manuscript writing and revising. BFY experiment performance, manuscript editing. ZKH experiment guide, manuscript revising. JQZ, YZ, TTX experiment performance.

#### Compliance with Ethical Standards

**Conflict of interest** There is no conflict of interest and all authors have read and approved the final manuscript.

## References

- Abdel Moneim AE (2016) *Indigofera oblongifolia* prevents lead acetate-induced hepatotoxicity, oxidative stress, fibrosis and apoptosis in rats. PLoS ONE 11:e0158965. doi:10.1371/journal.pone.0158965
- Abdou HM, Hassan MA (2014) Protective role of omega-3 polyunsaturated fatty acid against lead acetate-induced toxicity in liver and kidney of female rats. BioMed Res Int 2014:435857 doi:10.1155/2014/435857
- Aebi HE (1983) Catalase. In Bergmeyer HU (eds) Methods of Enzymatic Analysis, vol 3. Verlag Chemie, Weinheim, p 273–285
- Agency for Toxic Substances and Disease Registry (ATSDR) (2005) Toxicological profile for lead. US Department of Health and Human Services, Public Health Service, Atlanta, p 1–577
- Ali B, Song WJ, Hu WZ, Luo XN, Gill RA, Wang J, Zhou WJ (2014) Hydrogen sulfide alleviates lead-induced photosynthetic and ultrastructural changes in oilseed rape. Ecotoxicol Environ Saf 102:25–33. doi:10.1016/j.ecoenv.2014.01.013
- Al-Magableh MR, Kemp-Harper BK, Ng HH, Miller AA, Hart JL (2014) Hydrogen sulfide protects endothelial nitric oxide function under conditions of acute oxidative stress in vitro. Naunyn-Schmiedeberg's Arch Pharmacol 387:67–74 doi:10.1007/s00210-013-0920-x
- Beauchamp C, Fridovich I (1971) Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. Anal Biochem 44:276–287
- Bharwana SA, Ali S, Farooq MA, Ali B, Iqbal N, Abbas F, Ahmad MS (2014) Hydrogen sulfide ameliorates lead-induced morphological, photosynthetic, oxidative damages and biochemical changes in cotton. Environ Sci Pollut Res Int 21:717–731. doi:10.1007/s11356-013-1920-6
- Bian JS, Olson KR, Zhu YC (2016) Hydrogen sulfide: biogenesis, physiology, and pathology. Oxid Med Cell Longev 2016:6549625 doi:10.1155/2016/6549625
- Chen J, Zhu C, Li LP, Sun ZY, Pan XB (2007) Effects of exogenous salicylic acid on growth and H<sub>2</sub>O<sub>2</sub>-metabolizing enzymes in rice seedlings under lead stress. J Environ Sci 19:44–49
- Chen XY et al (2009) Endogenous hydrogen peroxide plays a positive role in the upregulation of heme oxygenase and acclimation to oxidative stress in wheat seedling leaves. J Integr Plant Biol 51:951–960. doi:10.1111/j.1744-7909.2009.00869.x
- Chen J et al (2011) Hydrogen sulphide enhances photosynthesis through promoting chloroplast biogenesis, photosynthetic enzyme expression, and thiol redox modification in *Spinacia oleracea* seedlings. J Exp Bot 62:4481–4493. doi:10.1093/jxb/err145
- Chen Z, Pan Y, Wang S, Ding Y, Yang W, Zhu C (2012) Overexpression of a protein disulfide isomerase-like protein from *Methanothermobacter thermoautotrophicum* enhances mercury tolerance in transgenic rice. Plant Sci 197:10–20. doi:10.1016/j.plantsci.2012.08.005
- Chen J et al (2013a) Hydrogen sulfide alleviates aluminum toxicity in barley seedlings. Plant Soil 362:301–318. doi:10.1007/s11104-012-1275-7
- Chen Z, Pan YH, An LY, Yang WJ, Xu LG, Zhu C (2013b) Heterologous expression of a halophilic archaeon manganese superoxide dismutase enhances salt tolerance in transgenic rice. Russ J Plant Physiol + 60:359–366 doi:10.1134/S1021443713030059
- Chen Z, Zhang L, Zhu C (2015) Exogenous nitric oxide mediates alleviation of mercury toxicity by promoting auxin transport in roots or preventing oxidative stress in leaves of rice seedlings. Acta Physiol Plant. doi:10.1007/s11738-015-1931-7
- Christou A, Manganaris GA, Papadopoulos I, Fotopoulos V (2013) Hydrogen sulfide induces systemic tolerance to salinity and non-ionic osmotic stress in strawberry plants through modification of reactive species biosynthesis and transcriptional regulation of multiple defence pathways. J Exp Bot 64:1953–1966. doi:10.1093/jxb/ert055
- Dewanjee S et al (2015) Water spinach, *Ipomoea aquatica* (Convolvulaceae), ameliorates lead toxicity by inhibiting oxidative stress and apoptosis. PLoS ONE 10:e0139831. doi:10.1371/journal.pone.0139831
- Dua TK et al (2016) Cytoprotective and antioxidant effects of an edible herb, *Enhydra fluctuans* Lour. (Asteraceae), against experimentally induced lead acetate intoxication. PLoS ONE 11:e0148757. doi:10.1371/journal.pone.0148757
- Fang Y, Hu XH, Jia ZG, Xu MH, Guo ZY, Gao FH (2012) Tiron protects against UVB-induced senescence-like characteristics in human dermal fibroblasts by the inhibition of superoxide anion production and glutathione depletion. Australas J Dermatol 53:172–180. doi:10.1111/j.1440-0960.2012.00912.x
- Fang T, Cao ZY, Li JL, Shen WB, Huang LQ (2014) Auxin-induced hydrogen sulfide generation is involved in lateral root formation in tomato. Plant Physiol Biochem 76:44–51 doi:10.1016/j.plaphy.2013.12.024
- Gupta S, Gupta M (2015) Alleviation of selenium toxicity in *Brassica juncea* L.: salicylic acid-mediated modulation in toxicity indicators, stress modulators, and sulfur-related gene transcripts. Protoplasma. doi:10.1007/s00709-015-0908-0

- Han YH, Moon HJ, You BR, Kim SZ, Kim SH, Park WH (2009) The effects of buthionine sulfoximine, diethyldithiocarbamate or 3-amino-1,2,4-triazole on propyl gallate-treated HeLa cells in relation to cell growth, reactive oxygen species and glutathione. *Int J Mol Med* 24:261–268
- Hancock JT, Whiteman M (2016) Hydrogen sulfide signaling: interactions with nitric oxide and reactive oxygen species. *Ann N Y Acad Sci* 1365:5–14. doi:[10.1111/nyas.12733](https://doi.org/10.1111/nyas.12733)
- Hoagland DR, Arnon DI (1950) The water-culture method for growing plants without soil. *Calif Agric Exp Stn Circ* 347:1–32
- Hu LY et al (2012) Hydrogen sulfide prolongs postharvest shelf life of strawberry and plays an antioxidative role in fruits. *J Agric Food Chem* 60:8684–8693 doi:[10.1021/jf300728h](https://doi.org/10.1021/jf300728h)
- Hu ZR, Xie Y, Jin GF, Fu JM, Li HY (2015) Growth responses of two tall fescue cultivars to Pb stress and their metal accumulation characteristics. *Ecotoxicology* 24:563–572. doi:[10.1007/s10646-014-1404-6](https://doi.org/10.1007/s10646-014-1404-6)

- Mishra SKSD, Singhal GS (1991) Interrelationship between salt and light stress on the primary process of photosynthesis. *J Plant Physiol* 138:92–96
- Mittler R (2002) Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci* 7:405–410. doi:10.1016/S1360-1385(02)02312-9
- Mostofa MG, Rahman A, Ansary MM, Watanabe A, Fujita M, Tran LS (2015) Hydrogen sulfide modulates cadmium-induced physiological and biochemical responses to alleviate cadmium toxicity in rice. *Sci Rep* 5:14078. doi:10.1038/srep14078
- Murali Achary VM, Panda BB (2010) Aluminium-induced DNA damage and adaptive response to genotoxic stress in plant cells are mediated through reactive oxygen intermediates. *Mutagenesis* 25:201–209. doi:10.1093/mutage/gep063
- Nakano Y, Asada K (1981) Hydrogen peroxide is scavenged by ascorbate-specific peroxidase in spinach chloroplasts. *Plant Cell Physiol* 22:867–880
- Olson KR (2012) A practical look at the chemistry and biology of hydrogen sulfide. *Antioxid Redox Sign* 17:32–44 doi:10.1089/ars.2011.4401
- Rao MV, Paliyath G, Ormrod DP, Murr DP, Watkins CB (1997) Influence of salicylic acid on H<sub>2</sub>O<sub>2</sub> production, oxidative stress, and H<sub>2</sub>O<sub>2</sub>-metabolizing enzymes. Salicylic acid-mediated oxidative damage requires H<sub>2</sub>O<sub>2</sub>. *Plant Physiol* 115:137–149
- Rao F, Zhai Y, Sun F (2016) Punicalagin mollifies lead acetate-induced oxidative imbalance in male reproductive system. *Int J Mol Sci* 17:1269. doi:10.3390/ijms17081269
- Reis GS, de Almeida AA, de Almeida NM, de Castro AV, Mangabeira PA, Pirovani CP (2015) Molecular, biochemical and ultrastructural changes induced by Pb toxicity in seedlings of *Theobroma cacao* L. *PLoS ONE* 10:e0129696. doi:10.1371/journal.pone.0129696
- Saxena I, Shekhawat GS (2013) Nitric oxide (NO) in alleviation of heavy metal induced phytotoxicity and its role in protein nitration. *Nitric Oxide* 32:13–20 doi:10.1016/j.niox.2013.03.004
- Schaedle M, Bassham JA (1977) Chloroplast glutathione reductase. *Plant Physiol* 59:1011–1012. doi:10.1104/pp.59.5.1011
- Sharma P, Dubey RS (2005) Lead toxicity in plants. *Braz J Plant Physiol* 17:35–52
- Sharma P, Mishra KP (2006) Aluminum-induced maternal and developmental toxicity and oxidative stress in rat brain: response to combined administration of Tiron and glutathione. *Reprod Toxicol* 21:313–321. doi:10.1016/j.reprotox.2005.06.004
- Shi H, Ye T, Chan Z (2013) Exogenous application of hydrogen sulfide donor sodium hydrosulfide enhanced multiple abiotic stress tolerance in bermudagrass (*Cynodon dactylon* (L.) Pers.). *Plant Physiol Biochem* 71:226–234 doi:10.1016/j.plaphy.2013.07.021
- Shi HT, Ye TT, Chan ZL (2014) Nitric oxide-activated hydrogen sulfide is essential for cadmium stress response in bermudagrass (*Cynodon dactylon* (L.) Pers.). *Plant Physiol Biochem* 74:99–107 doi:10.1016/j.plaphy.2013.11.001
- Sidhu GP, Singh HP, Batish DR, Kohli RK (2016) Effect of lead on oxidative status, antioxidative response and metal accumulation in *Coronopus didymus*. *Plant Physiol* 105:290–296 doi:10.1016/j.plaphy.2016.05.019
- Singh HP, Kaur S, Batish DR, Sharma VP, Sharma N, Kohli RK (2009) Nitric oxide alleviates arsenic toxicity by reducing oxidative damage in the roots of *Oryza sativa* (rice). *Nitric Oxide* 20:289–297. doi:10.1016/j.niox.2009.02.004
- Sun J, Wang RG, Zhang X, Yu YC, Zhao R, Li ZY, Chen SL (2013) Hydrogen sulfide alleviates cadmium toxicity through regulations of cadmium transport across the plasma and vacuolar membranes in *Populus euphratica* cells. *Plant Physiol Biochem* 65:67–74 doi:10.1016/j.plaphy.2013.01.003
- Verma S, Dubey RS (2003) Lead toxicity induces lipid peroxidation and alters the activities of antioxidant enzymes in growing rice plants. *Plant Sci* 164:645–655. doi:10.1016/S0168-9452(03)00022-0
- Wang FJ, Zeng B, Sun ZX, Zhu C (2009) Relationship between proline and Hg<sup>2+</sup>-induced oxidative stress in a tolerant rice mutant. *Arch Environ Contam Toxicol* 56:723–731. doi:10.1007/s00244-008-9226-2
- Wang BL, Shi L, Li YX, Zhang WH (2010) Boron toxicity is alleviated by hydrogen sulfide in cucumber (*Cucumis sativus* L.) seedlings. *Planta* 231:1301–1309. doi:10.1007/s00425-010-1134-9
- Wang YQ, Li L, Cui WT, Xu S, Shen WB, Wang R (2012) Hydrogen sulfide enhances alfalfa (*Medicago sativa*) tolerance against salinity during seed germination by nitric oxide pathway. *Plant Soil* 351:107–119. doi:10.1007/s11104-011-0936-2
- Wang Y et al (2015b) Metabolomic analysis with GC-MS to reveal potential metabolites and biological pathways involved in Pb & Cd stress response of radish roots. *Sci Rep* 5:18296 doi:10.1038/srep18296
- Wierzbicka M, Obidzińska J (1998) The effect of lead on seed imbibition and germination in different plant species. *Plant Sci* 137:155–171. doi:10.1016/S0168-9452(98)00138-1
- Wu G, Wan F, Fu H, Li N, Gao H (2015) A matter of timing: contrasting effects of hydrogen sulfide on oxidative stress response in *Shewanella oneidensis*. *J Bacteriol* 197:3563–3572. doi:10.1128/JB.00603-15
- Xie ZZ, Liu Y, Bian JS (2016) Hydrogen sulfide and cellular redox homeostasis. *Oxid Med Cell Longev*. doi:10.1155/2016/6043038
- Yang N, Zhou FR, Wang JX (2016) Eco-toxicological effects of two kinds of lead compounds on forest tree seed in alkaline soil. *Environ Monit Assess* 188:201. doi:10.1007/s10661-016-5198-2
- Zhang H, Hu LY, Hu KD, He YD, Wang SH, Luo JP (2008) Hydrogen sulfide promotes wheat seed germination and alleviates oxidative damage against copper stress. *J Integr Plant Biol* 50:1518–1529. doi:10.1111/j.1744-7909.2008.00769.x
- Zhang H, Hu LY, Li P, Hu KD, Jiang CX, Luo JP (2010) Hydrogen sulfide alleviated chromium toxicity in wheat. *Biol Plant* 54:743–747. doi:10.1007/s10535-010-0133-9
- Zhang H et al (2011) Hydrogen sulfide acts as a regulator of flower senescence in plants. *Postharvest Biol Technol* 60:251–257 doi:10.1016/j.postharvbio.2011.01.006
- Zhao L, Xiong J, Li LP, Zhu C (2009) Low concentration of exogenous abscisic acid increases lead tolerance in rice seedlings. *Biol Plant* 53:728–732. doi:10.1007/s10535-009-0132-x